

SCALAR CASIMIR EFFECT WITH NON-LOCAL BOUNDARY CONDITIONS

Aram Saharian,^{1,2,3} Giampiero Esposito^{4,5}

¹*Department of Physics, Yerevan State University,*

1 Alex Manoogian Street, 375049 Yerevan, Armenia

²*Abdus Salam International Centre for Theoretical Physics, 34014 Trieste, Italy*

³*Departamento de Física-CCEN, Universidade Federal da Paraíba,*

58.059-970, J. Pessoa, PB C. Postal 5.008, Brazil

⁴*Istituto Nazionale di Fisica Nucleare, Sezione di Napoli,*

Complesso Universitario di Monte S. Angelo,

Via Cintia, Edificio N', 80126 Naples, Italy

⁵*Dipartimento di Scienze Fisiche, Complesso Universitario di Monte S. Angelo,*

Via Cintia, Edificio N', 80126 Naples, Italy

Abstract

Non-local boundary conditions have been considered in theoretical high-energy physics with emphasis on one-loop quantum cosmology, one-loop conformal anomalies, Bose–Einstein condensation models and spectral branes. We have therefore studied the Wightman function, the vacuum expectation value of the field square and the energy-momentum tensor for a massive scalar field satisfying non-local boundary conditions on a single and two parallel plates. Interestingly, we find that suitable choices of the kernel in the non-local boundary conditions lead to forces acting on the plates that can be repulsive for intermediate distances.

In our analysis of the Casimir effect for scalar fields [1], motivated by the work in Refs. [2, 3], we have considered the geometry of two parallel plates with non-local boundary conditions

$$n_{(j)}^\nu \partial_\nu \varphi(x^\mu) + \int d\mathbf{x}'_\parallel f_j(|\mathbf{x}_\parallel - \mathbf{x}'_\parallel|) \varphi(x'^\mu) = 0, \quad x = a_j, \quad (1)$$

where we use rectangular coordinates $x^\mu = (t, x^\perp, \mathbf{x}_\parallel)$, with $\mathbf{x}_\parallel = (x^2, \dots, x^D)$, and $n_{(j)}^\nu$ is the inward-pointing unit normal to the boundary at $x = a_j$. For the region between the plates the corresponding eigenvalues are solutions of the equation [1]

$$(z^2 - c_1 c_2) \sin z + (c_1 + c_2) z \cos z = 0, \quad (2)$$

where the coefficients c_j are determined by the Fourier transforms F_j of the kernel functions $f_j(x_\parallel)$ in the boundary conditions, i.e.

$$c_j \equiv (-1)^{j-1} a F_j(k_\parallel) \equiv (-1)^{j-1} a \int d\mathbf{x}_\parallel f_j(|\mathbf{x}_\parallel|) e^{i\mathbf{k}_\parallel \cdot \mathbf{x}_\parallel}. \quad (3)$$

The non-local boundary conditions (1) state that the normal derivative at a given point depends on the values of the field at other points on the boundary. The properties of the boundary are expressed by the kernel function f_j . In a sense, this setting is similar to that in electrodynamics for the spatial dispersion of the dielectric function ε , where ε depends on the wave vector by virtue of spatial dispersion. Similarly, our non-local boundary conditions engender dependence of the coefficient F_j in the eigenfunctions on the wave vector $\mathbf{k}_\parallel$.

The evaluation of the corresponding Wightman function is based on a variant of the generalized Abel–Plana summation formula below [1]:

$$\begin{aligned} \sum_{z=\lambda_n, iy_l} \frac{h(z)}{1 + \cos(z + 2\alpha_1) \sin z/z} &= -\frac{1}{2} \frac{h(0)}{1 - c_1^{-1} - c_2^{-1}} + \frac{1}{\pi} \int_0^\infty dz h(z) \\ &+ \frac{i}{\pi} \int_0^\infty dt \frac{h(te^{\pi i/2}) - h(te^{-\pi i/2})}{\frac{(t-c_1)(t-c_2)}{(t+c_1)(t+c_2)} e^{2t} - 1} \\ &- \frac{\pi \theta(c_j)}{2c_j} [g_j(c_j e^{\pi i/2}) + g_j(c_j e^{-\pi i/2})], \end{aligned} \quad (4)$$

where $g_j \equiv (z^2 + c_j^2)h(z)$. The application of this formula has made it possible for us to extract from the VEVs the parts resulting from the single plate and to present the part induced from the second plate in terms of integrals exponentially convergent for points away

from the boundary. The Wightman function turns out to be given by [1]

$$\begin{aligned}
\langle 0 | \varphi(x^\mu) \varphi(x'^\mu) | 0 \rangle &= \langle 0_S | \varphi(x^\mu) \varphi(x'^\mu) | 0_S \rangle_j + \frac{4}{(2\pi)^D} \int d\mathbf{k}_\parallel e^{i\mathbf{k}_\parallel \cdot (\mathbf{x}_\parallel - \mathbf{x}'_\parallel)} \\
&\times \int_{a\sqrt{k_\parallel^2 + m^2}}^\infty dt \frac{\cosh(tx_j + \tilde{\alpha}_j) \cosh(tx'_j + \tilde{\alpha}_j)}{\frac{(t-c_1)(t-c_2)}{(t+c_1)(t+c_2)} e^{2t} - 1} \\
&\times \frac{\cosh \left[(t-t') \sqrt{t^2/a^2 - k_\parallel^2 - m^2} \right]}{\sqrt{t^2 - k_\parallel^2 a^2 - m^2 a^2}}, \tag{5}
\end{aligned}$$

having defined $\tilde{\alpha} \equiv \frac{1}{2} \log((t - c_j)/(t + c_j))$.

Moreover, the vacuum stress in the direction orthogonal to the plates is uniform. This stress determines the vacuum forces acting on the plates, and the corresponding effective pressure reads as [1]

$$\begin{aligned}
p &= -\langle 0 | T_1^1 | 0 \rangle = -\frac{2S_{D-1}}{(2\pi)^D} \int_0^\infty du u^{D-2} \int_{\sqrt{u^2 + m^2}}^\infty \frac{t^2 dt}{\sqrt{t^2 - u^2 - m^2}} \\
&\times \left[\frac{(t - F_1(u))(t + F_2(u))}{(t + F_1(u))(t - F_2(u))} e^{2at} - 1 \right]^{-1}. \tag{6}
\end{aligned}$$

We have evaluated numerically the vacuum forces acting on the plates in the case of the kernel functions [1]

$$f_j(x) \equiv f_{0j} e^{-\eta_j x}. \tag{7}$$

The corresponding Fourier transforms $F_j(k_\parallel)$ are given by the formulae

$$F_j(k_\parallel) = \frac{\eta_j F_1^{(j)}}{(1 + k_\parallel^2/\eta_j^2)^{D/2}}, \tag{8}$$

where the parameters $F_1^{(j)}$ are defined by

$$F_1^{(j)} \equiv 2^{D-1} \pi^{\frac{D}{2}-1} \Gamma(D/2) \frac{f_{0j}}{\eta_j^D}. \tag{9}$$

We find that, for the values $F_1^{(1)} \lesssim -1.08$, the vacuum pressure is negative for all interplate distances and the corresponding vacuum forces are attractive. For the values $F_1^{(1)} > -1.08$ there are two values of the distance between the plates for which the vacuum forces vanish. These values correspond to equilibrium positions of the plates. Moreover, for values of the distance in the region between these positions the vacuum forces acting on plates are repulsive. Thus, the left equilibrium position is unstable and the right one is locally stable [1].

It might be interesting to investigate the relation, if any, with the findings in Ref. [4], where the authors obtain a repulsive Casimir force among parallel plates under the assumption of a suitable ultraviolet cut-off such that the regularized zero-point energy of the vacuum can be the source of non-vanishing cosmological constant driving the acceleration of the Universe.

Acknowledgments

The work of A. Saharian has been supported by the INFN, by ANSEF Grant No. 05-PS-hep-th-89-70, and in part by the Armenian Ministry of Education and Science, Grant No. 0124. The work of G. Esposito has been partially supported by PRIN *SINTESI*.

-
- [1] A.A. Saharian and G. Esposito, hep-th/0512041.
 - [2] M. Schröder, Rep. Math. Phys. **27**, 259 (1989).
 - [3] G. Esposito, Class. Quantum Grav. **16**, 1113 (1999).
 - [4] G. Mahajan, S. Sarkar, and T. Padmanabhan, astro-ph/0604265.